Cellphone Radio Frequency Radiation Studies

Cellphones are used by 95% of American adults.¹ Given the large number of users, any harmful effects associated with cellphone use could be a significant public health concern.

Cellphones use radio frequency radiation (RFR) to transmit signals. The U.S. Food and Drug Administration (FDA) nominated RFR for study by the National Toxicology Program (NTP), due to widespread human exposure and limited information about the potential health effects of long-term cellphone use.

What did NTP study?

NTP conducted toxicology studies in rats and mice to help clarify potential health hazards, including cancer risk, from exposure to RFR used in 2G and 3G cellphones. 2G and 3G networks were standard when the studies were designed and are still used for phone calls and texting.

The \$30 million NTP studies took more than 10 years to complete and are the most comprehensive assessment, to date, of health effects in animals exposed to RFR. The results will help guide other studies of newer technologies.





What did the studies find?

NTP studies found that exposure to high levels of RFR, like that used in 2G and 3G cellphones, was associated with:

- Clear evidence of tumors in the hearts of male rats. The tumors were malignant schwannomas.
- Some evidence of tumors in the brains of male rats. The tumors were malignant gliomas.
- Some evidence of tumors in the adrenal glands of male rats. The tumors were pheochromocytomas.

For female rats, and male and female mice, it was unclear, also known as equivocal, whether cancers observed in the studies were associated with exposure to RFR.

The conclusions were based on NTP's four categories of evidence that a substance may cause cancer:

- Clear evidence (highest)
- Some evidence
- Equivocal evidence
- No evidence (lowest)



PO Box 12233 • Research Triangle Park, NC 27709 Phone: 919-541-3345 • https://ntp.niehs.nih.gov



Do the rat and mouse findings apply to humans?

The findings in animals cannot be directly applied to humans for two key reasons:

- The exposure levels and durations were greater than what people may receive from cellphones.
- The rats and mice received RFR across their whole bodies, which is different from the more localized exposures humans may receive, like from a cellphone in their pocket or next to their head.

However, the studies question the long-held assumption that radio frequency radiation is of no concern as long as the energy level is low and does not significantly heat the tissues.

Did NTP find health effects other than cancer?

NTP found lower body weights among newborn rats and their mothers, especially when exposed to high levels of RFR during pregnancy and lactation, yet these animals grew to normal size. They also found that RFR exposure was associated with an increase in DNA damage. DNA damage, if not repaired, can potentially lead to tumors. Specifically, they found RFR exposure was linked with significant increases in DNA damage in the frontal cortex of the brain in male mice, the blood cells of female mice, and the hippocampus of male rats.

What factors contributed to the NTP conclusions?

In addition to seeing tumors in the male rats with higher exposures to RFR, NTP scientists also observed other changes in the hearts of exposed male and female rats that supported their conclusions.

The evidence for tumors in the brain and adrenal glands was not as strong as what NTP scientists saw in the heart. However, the type of brain cancer observed is similar to a type of brain tumor linked to heavy cellphone use in some human studies.³

Still, the effects observed in the animals were relatively rare. NTP scientists are not sure why male rats appear to be at greater risk for developing tumors compared to female rats.

How do the RFR levels used in the studies compare to typical human exposures?


The lowest exposure level used in the studies was equal to the maximum local tissue exposure currently allowed for cellphone users. This power level rarely occurs with typical cellphone use. The highest exposure level in the studies was four times higher than the maximum power level permitted for local tissues.



Why did NTP expose the animals' whole bodies to RFR?

Although many previous studies focused on exposure to the brain, NTP scientists wanted to make sure that they were considering health effects to the whole body, especially since many people do not hold their phones next to their head much of time.

Non-lonizing lonizing Low Frequency High Frequency



RFR used in cellphones is at the lower frequency and lower energy end of the electromagnetic spectrum.



What is the difference between electric and magnetic fields and RFR?

RFR is a subcategory of electric and magnetic fields (EMF), which are the invisible waves of force that surround any electrical device. The different types of EMF are distinguished by their frequencies. RFR is a form of low frequency, non-ionizing radiation that was thought to be relatively harmless.

How were the studies conducted?

Rats and mice were exposed to RFR in special chambers for up to two years, or most of their natural lives. NTP scientists looked for a range of cancers and noncancer health effects.

Exposure to RFR began in the womb for rats and at 5-6 weeks old for mice. The RFR exposure was intermittent, 10 minutes on and 10 minutes off, totaling about nine hours each day. The RFR levels ranged from 1.5 to 6 watts per kilogram of body weight in rats, and 2.5 to 10 watts per kilogram in mice.

The chambers were shielded rooms with a transmitting antenna that radiated RFR fields, plus rotating stirrers that generated a uniform field.^{4,5} Pilot studies established field strengths that did not raise animal body temperatures excessively.⁶

The rats and mice were exposed to whole body RFR at frequencies of 900 and 1900 megahertz, respectively, from two technologies – Code Division Multiple Access (CDMA) and Global System for Mobile Communications (GSM).

NTP and RFR experts from the National Institute of Standards and Technology (NIST) and the IT'IS Foundation designed and built the chambers specifically for these studies.



Chambers at the IIT Research Institute in Chicago where the studies took place

What is the difference between CDMA and GSM?

CDMA and GSM are two common ways of transmitting cellphone signals in the U.S. and Europe. There are substantial differences in signal structure that may result in different RFR exposures, so NTP wanted to expose the animals to both modulations.

In March 2018, a panel of external scientific experts thoroughly reviewed the draft NTP technical reports at a public meeting. The final conclusions represent the consensus between NTP and the panel.

How do the NTP studies relate to 4G, 5G, or Wi-Fi?

NTP studies of RFR used in 2G and 3G cellphones do not apply to 4G or 5G technologies. These newer technologies use different methods of signal modulation than NTP used in the studies. The NTP studies also did not investigate frequencies and modulations used for Wi-Fi.

What were the studies' strengths?

NTP was able to control exactly how much RFR the animals received — something that is not possible when studying human cellphone use.

Were there any surprise findings?

NTP found longer lifespans among the exposed male rats. This may be explained by an observed decrease in chronic kidney problems that are often the cause of death in older rats.

What will NTP do with the results of the studies?

NTP will provide these studies to the FDA and Federal Communications Commission. The agencies will review the information as they continue to monitor new research on the potential health effects of RFR.

Final reports and data tables are available on the NTP website at https://ntp.niehs.nih.gov/go/cellphone.



What are NTP's future plans for studying cellphone RFR and 5G wireless technology?

5G is the emergent technology that will eventually overtake the existing technology. In the meantime, people will continue to be exposed to RFR in the 700–2700 MHz range. As the 5G network is implemented, some of the signals used by the 5G network will use the same lower frequencies used by the older technology, but the 5G network will also use higher frequencies—up to 60,000 MHz—thereby exposing people to a much broader spectrum of frequencies. The higher frequencies, known as millimeter waves, can rapidly transmit enormous amounts of data with increased network capacity compared with current technologies. Millimeter waves do not travel as far and do not penetrate the body as deeply as do the wavelengths from the lower frequencies. Millimeter waves are likely to penetrate no deeper than the skin, whereas the lower frequencies have been shown to penetrate at least 3 to 4 inches into the human body.

NTP is currently evaluating the existing literature on the higher frequencies intended for use in the 5G network, and is working to better understand the biological basis for the cancer findings reported in earlier studies on RFR with 2G and 3G technologies. Additionally, work is ongoing to develop smaller RFR exposure chambers for additional short-term studies that will take weeks and months to complete rather than years. The exposure system is also being designed to have the capability to conduct studies with various RFR frequencies and modulations to keep up with the changing technologies in the telecommunications industry.

Where can I go for more information?

For more information on what federal agencies are doing to determine whether RFR used in cellphones may affect human health, visit the following websites:

National Toxicology Program

https://ntp.niehs.nih.gov/go/cellphone

National Cancer Institute

www.cancer.gov/about-cancer/causesprevention/risk/radiation/cell-phones-factsheet

U.S. Food and Drug Administration https://go.usa.gov/B5tx

Federal Communications Commission www.fcc.gov/consumers/guides/wireless-devices-and-health-concerns

The National Toxicology Program is an interagency program headquartered at the **National Institute of Environmental Health Sciences** that tests and evaluates chemicals in our environment.

For more information on NTP, go to https://ntp.niehs.nih.gov.

¹ Pew Research Center. 2018. Mobile Fact Sheet. Available: www.pewinternet.org/fact-sheet/mobile [accessed 24 October 2018].

² FDA (U.S. Food and Drug Administration). 2017. Reducing Exposure: Hands-free Kits and Other Accessories. Available: https://www.fda.gov/Radiation-EmittingProducts/RadiationEmittingProductsandProcedures/HomeBusinessandEntertainment/CellPhones/ucm116293.htm [accessed 30 October 2018].

³ IARC (International Agency for Research on Cancer). 2013. Non-lonizing Radiation, Part 2: Radiofrequency Electromagnetic Fields. IARC Monogr Eval Carcinog Risks Hum 102:1-481.

⁴ Capstick MH, Kuehn S, Berdinas-Torres V, Gong Y, Wilson PF, Ladbury JM, Koepke G, McCormick DL, Gauger J, Melnick RL, Kuster N. 2017. A radio frequency radiation exposure system for rodents based on reverberation chambers. IEEE T Electromagn C 59(4):1041-1052.

⁵ Gong Y, Capstick MH, Kuehn S, Wilson PF, Ladbury JM, Koepke G, McCormick DL, Melnick RL, Kuster N. 2017. Life-time dosimetric assessment for mice and rats exposed in reverberation chambers for the two-year NTP cancer bioassay study on cell phone radiation. IEEE T Electromagn C 59(6):1798-1808.

⁶ Wyde ME, Horn TL, Capstick MH, Ladbury JM, Koepke G, Wilson PF, Kissling GE, Stout MD, Kuster N, Melnick RL, Gauger J, Bucher JR, McCormick DL. 2018. Effect of cell phone radiofrequency radiation on body temperature in rodents: Pilot studies of the National Toxicology Program's reverberation chamber exposure system. Bioelectromagnetics 39(3):190-199.